



cerexagri



A S E L E C T I V E H E R B I C I D E

For Controlling Certain Unwanted Aquatic Plants

ACTIVE INGREDIENT:	
Butoxyethyl Ester of 2,4-Dichlorophenoxyacetic Acid*	27.6%
OTHER INGREDIENTS:	72.4%
TOTAL	100.0%

*Isomer Specific AOAC Method,
 Equivalent to 2,4-Dichlorophenoxyacetic Acid 19.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID:

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

EPA Registration No. 228-378-4581

EPA Establishment No. 228-IL-1

Net Contents _____

Distributed By:

Cerexagri, Inc.

2000 Market Street, Philadelphia, PA 19103
 1 800-438-6071 • www.cerexagri.com



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not use in or near a greenhouse.

OXYGEN RATIO

Fish breathe oxygen in the water and a water-oxygen ratio must be maintained. Decaying weeds use up oxygen, but during the period when this product should be used, the weed mass is fairly sparse and the weed decomposition rate is slow enough so that the water oxygen ratio is not disturbed by treating the entire area at one time.

If treatments must be applied later in the season when the weed mass is dense and repeat treatments are needed, spread granules in lanes, leaving buffer strips which can then be treated when vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment.

Buffer lanes should be 50 to 100 feet wide. Treated lanes should be as wide as the buffer strips (See illustration below).



WATER pH

Best results are generally obtained if the water to be treated has a pH less than 8. A pH of 8 or higher may reduce weed control. If regrowth occurs within a period of 6 to 8 weeks, a second application may be needed.

PERMIT TO USE CHEMICALS IN WATER

In many states, permits are required to control weeds by chemical means in public water. If permits are required, they may be obtained from the Chief, Fish Division, State Department of Conservation or the State Department of Public Health.

GENERAL INFORMATION

This product is formulated on special heat treated attaclay granules that resist rapid decomposition in water, sink quickly to lake or pond bottoms and release the weed killing chemical into the critical root zone area.

This product is designed to selectively control the weeds listed on the label. While certain other weeds may be suppressed, control may be incomplete. Reduced control may occur in lakes where water replacement comes from bottom springs.

WHEN TO APPLY

For best results, spread this product in the Spring and early Summer, during the time weeds start to grow. If desired, this timing can be checked by sampling the lake bottom in areas heavily infested with weeds the year before.

If treatments are delayed until weeds form a dense mat or reach the surface, two treatments may be necessary. Make the second treatment when weeds show signs of recovery. Treatments made after September may be less effective depending upon water temperatures and weed growth.

Occasionally, a second application will be necessary if heavy regrowth occurs or weeds reinfest from untreated areas.

HOW TO APPLY

FOR LARGE AREAS: Use a fertilizer spreader or mechanical seeder such as the Gerber or Gandy or other equipment capable of uniformly applying this product. Before spreading any chemical, calibrate your method of application to be sure of spreading the proper amount. When using boats and power equipment, you must determine the proper combination of (1) boat speed (2) rate of delivery from the spreader, and (3) width of swath covered by the granules.

FOR SMALL AREAS (Around Docks or Isolated Patches of Weeds):

Use a portable spreader such as the Cyclone seeder or other equipment capable of uniformly applying this product. Estimate or measure out the area you want to treat. Weigh out the amount of material needed and spread this uniformly over the area. More uniform coverage is obtained by dividing the required amount in two and covering the area twice, applying the second half at right angles to the first.

Use the following formula to calibrate your spreader's delivery in pounds of this product per minute.

$$\frac{\text{Miles per hour} \times \text{spreader width} \times \text{pounds per acre}}{495}$$

Example: To apply 100 pounds of this product per acre using a spreader that covers a 20 foot swath from a boat traveling at 4 miles per hour, set the spreader to deliver 16 pounds of this product per minute.

$$\frac{4 \text{ mph} \times 20 \text{ feet} \times 100 \text{ lbs./A}}{495}$$

AMOUNTS TO USE

Rates of application vary with resistance of weed species to the chemical, density of weed mass at time of treatment, stage of growth, water depth, and rate of water flow through the treated area. Use the higher rate for dense weeds, when water is more than 8 feet deep and where there is a large volume turnover.

	POUNDS PER ACRE	POUNDS PER 2000 SQ. FT.
SUSCEPTIBLE WEEDS Water milfoil (<i>Myriophyllum</i> spp.) Water stargrass (<i>Heteranthera dubia</i>)	100 to 200	5
SLIGHTLY TO MODERATELY RESISTANT WEEDS Bladderwort (<i>Utricularia</i> spp.) White water lily (<i>Nymphaea</i> spp.) Yellow water lily or (Nuphar spp.) spatterdock* Water shield (<i>Brasenia</i> spp.) Water chestnut (<i>Trapa natans</i>) Coontail* (<i>Ceratophyllum demersum</i>)	150 to 200	7½ to 10

*Repeat treatments may be needed.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)**

CAUTION

Harmful if swallowed, absorbed through skin, or inhaled. Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. When handling this product, wear chemical resistant gloves. Wash thoroughly with soap and water after handling. When mixing, loading, or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow and temple protection. Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift or runoff may adversely affect fish and nontarget plants. Do not apply to water except as specified on this label. Do not contaminate water when disposing of equipment washwaters. Unless an approved assay indicates the 2,4-D concentration is 100 ppb (0.1 ppm) or less, or, only growing crops and noncrop areas labeled for direct treatment with 2,4-D will be affected, do not use water from treated areas for irrigating plants or mixing sprays for agricultural or ornamental plants.

Unless an approved assay indicates the 2,4-D concentration is 70 ppb (0.07 ppm) or less, do not use water from treated areas for potable water (drinking water).

Clean spreader equipment thoroughly before using it for any other purposes. Vapors from this product may injure susceptible plants.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal. It is recommended that a SARA Title III emergency response plan be created for storage facilities. Do not transport in the passenger compartment of any vehicle.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal or excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay away from smoke.

EMERGENCY TELEPHONE NUMBERS:

CHEMTREC: (800) 424-9300

MEDICAL: (303) 623-5716

Rocky Mountain Poison Control Center

WARRANTY AND DISCLAIMER

Cerexagri, Inc. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. CEREXAGRI MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL CEREXAGRI OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COST; OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

Cerexagri and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of Cerexagri.

Cerexagri, Inc. is a wholly-owned subsidiary of ATOFINA Chemicals, Inc.

Aqua-Kleen is a registered trademark of Nufarm, Inc.

5-F106S-01 A1 (1/01)

Made and Printed in U.S.A.