

Prescription Treatment[®] brand **AVERT[®]**

Pressurized Cockroach Bait Formula 1

KILLS: Cockroaches (including carbamate, organophosphate, organochlorine and pyrethroid resistant strains).

FOR USE IN: Garages, Homes, and the non-food/feed areas of Hospitals and Nursing Homes (non-occupied patient areas), Hotels, Motels, Transportation Equipment (Buses, Boats, Ships, Trains, and Planes[†]), Utilities, Warehouses, and Other Commercial and Industrial buildings.

[†] Do not use in aircraft cabins.

Not for use in USDA Inspected Meat and Poultry Plants

ACTIVE INGREDIENT:

Abamectin B1 [A mixture of avermectins containing 80% avermectin B1a (5-O-demethyl avermectin A1a) and 20% avermectin B1b (5-O-demethyl-25-de(1-methylpropyl)-25-(1-methylethyl) avermectin A1a)] 0.01%

INERT INGREDIENTS: 99.99%

Contains petroleum distillates.

EPA Reg. No. 499-322 EPA Est. No. 499-MO-1
KEEP OUT OF REACH OF CHILDREN
CAUTION

See side panel for additional precautionary statements

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

HOW TO USE WHITMIRE MICRO-GEN CRACK AND CREVICE[®] INJECTION SYSTEM: Whitmire Micro-Gen PT[®] Avert[®] Pressurized Cockroach Bait is for use only with the supplied actuator and injection tubes. PT[®] Avert[®] Pressurized Cockroach Bait is to be injected only into cracks and crevices or void space where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one half to one second of product. Concentrate treatment at insect activity sites. For light infestations, a minimum of 4 - 6 bait points is recommended per 100 sq. feet of treatment area. For heavy infestations, a minimum of 12 - 24 bait points is recommended per 100 sq. feet of treatment area. For inaccessible voids drill small holes and inject PT[®] Avert[®] Pressurized Cockroach Bait. Repeat as necessary.

Do not use in the food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food/feed is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but exclude areas where foods may be produced or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

Non-food/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Do not use in aircraft cabins.

CRAWLING INSECTS - Cockroaches (including carbamate, organophosphate, organochlorine and pyrethroid resistant strains): Inject the recommended dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, meter boxes and man holes. Repeat as necessary.

OUTDOOR USE: Use for control of cockroaches. Inject into cracks and crevices around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces and other areas where pests hide, such as tree holes and cracks in fences.

STORAGE & DISPOSAL

Do not puncture or incinerate. Do not contaminate water, food or feed by storage or disposal. Not for Storage in and around the home.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: This container may be recycled in the few but growing number of communities where aluminum aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label. (DO NOT PUNCTURE!). If recycling is not available, wrap the container and discard in the trash.

PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Give 1 or 2 glasses of water and immediately induce vomiting by gently touching back of throat with finger or administer syrup of Ipecac. Do not give anything by mouth to an unconscious person.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Seek medical attention if irritation persists.

IF ON EYES: Flush eyes with plenty of water. Get medical attention immediately.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. May be absorbed through the skin. Avoid breathing of spray mist and provide adequate ventilation of area being treated. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before smoking or eating. Avoid contamination of food, utensils, and food preparation surfaces. For spot treatment only. Do not use as a space spray. Do not allow children or pets in treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting. Do not spray directly into any electronic equipment such as radios, televisions, computers, etc. Do not spray where electrical short circuits might result, such as wall outlets, conduits, etc.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit the use of CFC propellants in aerosols.



A Prescription Treatment[®] brand insecticide from:
Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
<http://www.wmmg.com>

© 1998 Whitmire Micro-Gen Research Laboratories, Inc.

Revised 11/98