



TRIMEC CLASSIC BRAND BROADLEAF HERBICIDE

For Sale To and Use By Professional Turf Maintenance Personnel, Landscaping or Commercial Applicators Only.
Not For Sale To or Use By Homeowners.

Low-odor formulation!
Outstanding broadleaf weed control.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	25.93%
Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	13.85%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid	2.76%
INERT INGREDIENTS:	57.46%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

2.03 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 21.54%.
1.08 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 11.45%.
0.21 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.29%.
Isomer Specific by AOAC Method 6.D01-5
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN DANGER

See below for additional Precautionary Statements and Statement of Practical Treatment.

ONE GALLON COVERS 2 OR MORE ACRES

881/10-2000 AP021100
EPA REG. NO. 2217-543

Manufactured By
G pbi/gordon CORPORATION
An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin. Avoid breathing vapor or spray mist.

FOR TURF SITES ONLY

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

Clothing Requirement Statement: When mixing, loading, or applying this product, wear long-sleeved shirt, long pants, socks, shoes, chemical-resistant gloves and eye protection. It is recommended that safety glasses include front, brow and temple protection.

Personal Hygiene Statement: After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Engineering Control Statement:

Containers over 1 gallon and less than 5 gallons: Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

Containers of 5 gallons or more: Do not open-pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Statement of Practical Treatment

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

IF INHALED: Remove victim to fresh air and apply respiration if indicated.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. Do not contaminate domestic or irrigation waters. Do not apply when weather conditions favor drift from target area.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For metal containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or if allowed by other procedures approved by state and local authorities. **For plastic containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

USE PRECAUTIONS/BROADLEAF WEEDS IN ORNAMENTAL LAWNS AND TURF:

Do not apply this product through any type of irrigation system. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Use only lawn type sprayers. Avoid fine sprays; coarse sprays are less likely to drift. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other ornamental species. Do not apply to newly seeded grasses until well established. Do not

broadcast apply when air temperatures exceed 85°F; some injury may be expected with spot treatments when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application. After using this product, clean sprayer with soap or detergent and rinse thoroughly before using other pesticides in it. Care should be taken not to make applications where runoff could contaminate bodies of water or spray drift could carry the chemical to food crops or grazing lands where cattle, sheep, goats, swine or poultry would be exposed.

FOR USE ON RESIDENTIAL AND OTHER TURF SITES EXCLUDING SOD FARMS:

TRIMEC® Classic Brand Broadleaf Herbicide is intended to be applied by lawn care operators and landscape personnel for use in ornamental lawns and turf established in institutional, ornamental, and residential/domestic sites. Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses, and office buildings. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings. Finally, residential/domestic sites are defined as areas associated with the household or home life including but not limited to apartment complexes, condominiums and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

ORNAMENTAL LAWNS ESTABLISHED WITH COOL SEASON AND WARM SEASON TURFGRASSES:

Maximum control of weeds in cool season and warm season turfgrasses will be obtained from spring or early fall applications when weeds are actively growing. Do not use on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present.

Avoid spraying during long, excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours after application. Do not apply to newly seeded turf until after the second or third mowing. Grass seed can be sown 3 to 4 weeks after application at recommended rates.

The maximum application rate to turf is 1.0 lb. 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

General Application:

Cool Season Grasses, Common Bermudagrass, Bahiagrass and Zoysiagrass — Apply 3.25-4.0 pints of product per acre with spray volumes of 20-260 gallons per acre (gpa). Or, apply 1.2 to 1.5 fluid ounces of product in 0.5 to 6 gallons of water per 1,000 square feet. Use the higher application rate with the higher spray volumes. The best time to control clover is in the fall. When making a spring application to control clover, a 4 pint per acre rate is recommended.

Controlled Droplet Applicator (CDA):

Cool Season Grasses, Common Bermudagrass, Bahiagrass and Zoysiagrass — Add 2.5 pints of product to the Herbi container and fill with water. Spray contents over 33,000 square feet. Avoid overlapping between spray patterns.

Hybrid Bermudagrass — Apply 2.0-4.0 pints of product per acre with spray volumes of 20-260 gallons per acre (gpa). Or, apply 0.75 to 1.5 fluid ounces of product in 0.5 to 6 gallons of water per 1,000 square feet. Use the higher application rate with the higher spray volumes.

Certain hybrid bermudagrasses vary in tolerance to this product. Contact your local Extension Service Weed Control Specialist for turfgrass tolerance to this product.

Bentgrass, Centipedegrass and St. Augustinegrass — On closely mowed bentgrass (putting greens, bowling greens, and show lawns), apply a maximum rate of 1.0 fluid ounce of product per 1,500 square feet with spray volumes of 5.0 gallons per 1,500 square feet. Do not apply when grass is under stress from drought or hot temperatures. Use application equipment that is capable of distributing uniform droplets over the entire area to be sprayed. Slight yellowing of turf will disappear after about one week. Care should be taken to avoid overdosing bentgrass or injury may result.

Some injury can be expected to turf when used on centipedegrass or St. Augustinegrass lawns; spot spray weeds established in centipedegrass and St. Augustinegrass to avoid damage that may occur from a broadcast treatment. Do not apply this product to St. Augustinegrass during spring green-up which is the transition period between dormancy and active growth. Cultivars of St. Augustinegrass vary in tolerance to this product. Do not apply this product to 'Floritam' St. Augustinegrass.

BROADLEAF WEEDS

Aster, white heath & white prairie	Filaree, whitestem & redstem	Prostrate knotweed (*knotweed)
Bedstraw	Florida pusley	Puncturevine
Beggarweed, creeping	Ground ivy	Purple cudweed
Bindweed	Groundsel	Purslane
Black medic	Hawkweed	Ragweed
Broadleaf plantain	Healall	Red sorrel (*sheep sorrel)
Buckhorn plantain	Henbit	Shepherdspurse
Bull thistle	Innocence	Spotted spurge
Burclover	(Blue-eyed Mary)	Spurge
Burdock, common	Lambsquarters	Thistle
Buttercup, creeping	Lawn burweed	<i>Veronica</i>
Carpetweed	Lespedeza, common	(*corn speedwell)
Chickweed, common	Mallow, common	Virginia buttonweed
Chicory	Matchweed	White clover (*Dutch clover, honeysuckle clover, white trefoil & purplewort)
Cinquefoil	Mouseear chickweed	Wild carrot
Clover	Old world diamond flower	Wild garlic
Compassplant	<i>Oxalis</i> (*yellow woodsorrel & creeping woodsorrel)	Wild geranium
Curly dock	Parsley-piert	Wild lettuce
Dandelion	Pennsylvania smartweed	Wild mustard
Dayflower	Pennywort	Wild onion
Deadnettle	(*dollarweed)	Wild strawberry
Dock	Pepperweed	Wild violet
Dogfennel	Pigweed	Yarrow
English daisy	Pineappleweed	Yellow rocket
False dandelion (*spotted catsear & common catsear)	Plantain	
Field bindweed (*morningglory & creeping jenny)	Poison ivy	
Field oxeye-daisy (*creeping oxeye)	Poison oak	
*Synonyms		

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.